UPON ENROLMENT, AN ULTRACARE CASE MANAGER WILL:

- Partner with and remain dedicated to your patient throughout the treatment journey
- · Contact the patient or caregiver to review insurance coverage and support programs

## **Getting Started: Steps for Successful Enrolment in UltraCare**

Below are the steps for ensuring complete and timely enrolment in UltraCare so your patient can benefit fully from the program.

**OBTAIN PATIENT CONSENT**<sup>a</sup> The patient signature or verbal consent is required to allow third parties to share protected health information with Ultragenyx

2 SELECT PREFERRED PATIENT COMMUNICATION METHOD Ask your patient and/or caregiver about how they will prefer to communicate with their UltraCare Case Manager and the best time to contact them

**3** PRESCRIBER INFORMATION

ultragenyX UltraCare<sup>®</sup>

Provide contact details

4

**SPECIFY PRESCRIPTION FOR Pr DOJOLVI® (triheptanoin)** 

Provide a wet signature and date, which are necessary to process the prescription

alf the patient wants to opt out of the patient consent section, inform the UltraCare team on the phone or in writing by emailing ultracare@innomar-strategies.com.

## PATIENT CONSENT TO COLLECT, USE AND SHARE PERSONAL INFORMATION (PI) AND SIGNATURE

I understand that the UltraCare Program ("Program") is sponsored by Ultragenyx Pharmaceutical, Inc. ("Ultragenyx") and administered by Innomar on behalf of Ultragenyx. I understand that other service providers may be appointed by Ultragenyx to administer the Program from time to time. I authorize each of my physicians and pharmacists (including any specialty pharmacies and other healthcare providers), and each of my health insurers, to disclose my Pl, including but not limited to medical records, information related to my medical condition and treatment, financial, lab values, insurance coverage information, my name, address and telephone number to Ultragenyx and its agents, contractors, and assignees who will collect, use and disclose my Pl to manage and administer the Program, including to enrol me in and contact me about UltraCare Patient Services, provide case management through telephone or electronic communications to assist with adherence to my medication regimen, and work with third parties to provide community resources and referrals. I also authorize the collection, use and disclosure of information provided directly by me to the Program for legal obligations to report adverse drug events to health authorities and to monitor product complaints. I understand that Ultragenyx may contact me or my healthcare providers for additional information to fulfill its reporting obligations. I also understand that my Pl may be combined with the information of others who participate in the Program in order to generate aggregated data to improve the Program, to design and implement other patient programs and for research purposes including the identification of trends such as product utilization, adherence or outcomes.

I understand that Ultragenyx and its agents, contractors and assignees may store or process my PI outside of Canada (including in the United States), where local laws may require the disclosure of PI to government authorities under circumstances that are different than those that apply in Canada. I understand I may refuse to sign this consent, in which case I cannot be enrolled in the Program and understand that my treatment and eligibility for health benefits, including my access to therapy, will not be otherwise conditioned on my signing this consent. I understand that revoking this consent will not affect the ability to use and disclose PI received prior to receipt of notification that I wish to discontinue my participation in the Program. I understand I may revoke this consent at any time verbally or by writing to the address listed at the top of this form. Once consent has been revoked, I understand no additional PI will be collected. I understand that my PI will not be used or disclosed for any purposes, unless permitted by law, other than the purposes stated herein.

I understand that I may contact the Program at any time to update or access my PI, modify, express a privacy-related concern, or inquire about the privacy practices of the Program.

Patient Signature

Parent/Guardian Signature (if patient is a minor)

**IMPORTANT:** If healthcare provider is unable to obtain written consent from patient, please document when patient verbal consent was obtained. This will allow the program to continue with processing this enrolment. Written consent will be obtained by the program. Verbal consent should be obtained by a healthcare provider.

Patient consented verbally Date (DD/MM/YYYY) \_\_\_\_

Patient consent obtained by: Nam	e (Last, First)
----------------------------------	-----------------

\_ Title: MD RN Other (specify) \_

Signature

By providing my email address, I agree to receive, electronically, communications from Innomar acting on behalf of Ultragenyx Pharmaceutical, Inc. containing information and updates relating to my enrolment in the UltraCare Program. I understand that I may withdraw my consent to such communications at any time by providing notice to Innomar Strategies, Inc., c/o UltraCare Program, 2600 Alfred Nobel Blvd., Ville Saint-Laurent, QC H4S 0A9, or via email at ultracare@innomar-strategies.com.

You can report any suspected side effects associated with the use of health products to Health Canada at 1-866-234-2345 or http://www.hc-sc.gc.ca/dhp-mps/medeff/report-declaration/index-eng.php. You may also report side effects to Ultragenyx at 1-833-388-5872 (U-LTRA).

Toll-free Line: 1-833-388-5872 (U-LTRA) | Fax: 1-833-592-2273 (CARE) | http://www.ultracaresupport.ca | Email: Ultracare@innomar-strategies.com

\_\_\_\_\_ Date \_\_\_\_

## **Patient Enrolment Form**

PATIENT INFO						
		ose your preferred cont		PRESCRIBER INFORMATION	:	
, ,	ist Name			First Name		
	male 🗌 Male DOB (DD/N			Last Name		<u>.</u>
	nber			Street Address		
Street Address				City		
				Province	Postal Code	
Province		Postal Code	3			
Home Phone (_	) Wo	ork Phone ()				
Mobile Phone (	) E	Best Time to Contact		Fax ()		
Preferred Meth	od of Contact: 🗌 Home 🛽	Work Mobile	Email	Office Email		
Preferred Lang	uage: 🗌 English 📃 French	n Other		Office Contact Name/Title		
Email				Office Contact Phone ()		
Caregiver Nam	e (First and Last)			License #		
Relationship to	PatientCare	giver Phone ()	) [			
Has the patien	t previously been on MCT?	Yes What was the	dose?		No	
Confirmed diag LC-FAOD by (ch	neck one) acylcarnitii	nes on a newborn	lymphocytes) below	the lower limit of the normal mu	netic testing demonstr utation in a gene associ ain fatty acid oxidation	ated with long-
	ncy Type 🗌 CPTI 🔄 ( diagnosis of OTHER					
Initiate DOJOLVI over a period of For patients swit	2 to 3 weeks. Thing from another MCT produce	oximately 10% DCI divided i	c <u>t</u> nto at least 4 times pe	aloric intake (DCI), converted to mL, divid r day and increase to the recommended dosage of MCT divided into at least 4 tim	total daily dosage of up	to 35% DCI
Initiate DOJOLVI over a period of For patients swit Discontinue use dosage by appro <b>The total daily d</b> • Caloric value of	at a total daily dosage of appro 2 to 3 weeks. ching from another MCT produ- of MCT products before starting primately 5% DCI every 2 to 3 da <b>losage (mL) of DOJOLVI is dete</b> of DOJOLVI = 8.3 kcal/mL	n triglyceride (MCT) produ xximately 10% DCI divided i ct g DOJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin	ct nto at least 4 times pe /I at the last tolerated o of up to 35% DCI is acl	r day and increase to the recommended dosage of MCT divided into at least 4 tim nieved.	total daily dosage of up nes per day. Increase the <u>CI (kcal) x Target</u>	to 35% DCI total daily % <i>dose of DCI</i>
Initiate DOJOLVI over a period of For patients swift Discontinue use dosage by appro The total daily d • Caloric value c • Round the total	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting ximately 5% DCI every 2 to 3 da <b>losage (mL) of DOJOLVI is det</b> of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w al daily dosage into at least 4 ap	n triglyceride (MCT) produ xximately 10% DCI divided i DOJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin hole number	ct nto at least 4 times pe /I at the last tolerated o of up to 35% DCI is ach <b>ig calculation:</b>	r day and increase to the recommended dosage of MCT divided into at least 4 tin nieved. Patients D	total daily dosage of up nes per day. Increase the	to 35% DCI total daily % <i>dose of DCI</i>
Initiate DOJOLVI over a period of For patients swii Discontinue use dosage by appro The total daily d • Caloric value c • Round the tota • Divide the tota DOJOLVI Prescription	at a total daily dosage of appro 2 to 3 weeks. aching from another MCT produ- of MCT products before starting oximately 5% DCI every 2 to 3 da <b>losage (mL) of DOJOLVI is dete</b> of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w	n triglyceride (MCT) produ xximately 10% DCI divided i DOJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin hole number	ct nto at least 4 times pe /I at the last tolerated o of up to 35% DCI is ach ig calculation: al doses Initial mL	r day and increase to the recommended dosage of MCT divided into at least 4 tim nieved. $Total Daily Dose (mL) = \frac{Patients D}{2}$	total daily dosage of up nes per day. Increase the CI (	to 35% DCI total daily % <i>dose of DCI</i>
Initiate DOJOLVI over a period of For patients swift Discontinue use dosage by appro The total daily d • Caloric value c • Round the tot. • Divide the tot.	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting oximately 5% DCI every 2 to 3 da <b>losage (mL) of DOJOLVI is dete</b> of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w al daily dosage into at least 4 ap <b>linitial Total Daily Dose (mL)</b> Rounded to Nearest Whole Number	In triglyceride (MCT) produ iximately 10% DCI divided i g DOJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin hole number proximately equal individu ÷ Doses/Day =	ct nto at least 4 times pe /I at the last tolerated o of up to 35% DCI is ach ig calculation: al doses Initial mL	r day and increase to the recommended dosage of MCT divided into at least 4 tim nieved. <i>Total Daily Dose</i> (mL)= <i>Patients D</i> Increase bymL every until reaching targetmL d Use the Prescription Directions field below	total daily dosage of up nes per day. Increase the CI (	to 35% DCI total daily % <i>dose of DCI</i> //
Initiate DOJOLVI over a period of For patients swill Discontinue use dosage by appro The total daily d • Caloric value c • Round the tot. • Divide the tot. • DojOLVI Prescription Divide DOJOLVI Prescription Divide	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting oximately 5% DCI every 2 to 3 da <b>losage (mL) of DOJOLVI is dete</b> of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w al daily dosage into at least 4 ap <b>linitial Total Daily Dose (mL)</b> Rounded to Nearest Whole Number	In triglyceride (MCT) produ iximately 10% DCI divided i g DOJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin hole number proximately equal individu ÷ Doses/Day = (at least 4) mL) ÷	ct nto at least 4 times pe /I at the last tolerated o of up to 35% DCI is ach ig calculation: al doses Initial mL	r day and increase to the recommended dosage of MCT divided into at least 4 tim nieved. <i>Total Daily Dose</i> (mL)= <i>Patients D</i> Increase bymL every until reaching targetmL d Use the Prescription Directions field below	total daily dosage of up nes per day. Increase the CI (	to 35% DCI total daily % <i>dose of DCI</i> // Refills
Initiate DOJOLVI over a period of For patients swill Discontinue use dosage by appro The total daily d • Caloric value c • Round the tot. • Divide the tot. • Divide the tot. • Divide the tot. • DOJOLVI Prescription (Titration) Prescription Din DOJOLVI Prescription Din How Supplied: D Please see full Pl authorize PSP to	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting oximately 5% DCI every 2 to 3 da losage (mL) of DOJOLVI is detect of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w al daily dosage into at least 4 ap Initial Total Daily Dose (mL) Rounded to Nearest Whole Number Target Total Daily Dose ( Rounded to Nearest Whole Number DOJOLVI (triheptanoin) oral liquid roduct Monograph atwww.ultraget	In triglyceride (MCT) produ iximately 10% DCI divided i pDOJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin hole number proximately equal individu $\dot{+}$ Doses/Day = (at least 4) $\dot{+}$ (at is supplied in 500 mL bott genyx.com/canada/medici ward this prescription by fa	ct         nto at least 4 times per         /l at the last tolerated of up to 35% DCl is acl         ig calculation:         al doses         Initial mL per Dose	r day and increase to the recommended dosage of MCT divided into at least 4 tim nieved.  Total Daily Dose (mL)=  Patients D until reaching targetmL d Use the Prescription Directions field belo describe alternate desired dosing proto  Days Supply  Donograph-CANADA/ for complete dosa bivery, to the pharmacy chosen by the alternate dosine by the alternate do	total daily dosage of up hes per day. Increase the CI (kcal) x Target 8.3 kcal of DOJOLI day(s) Days Supply Refill age and administration ir	to 35% DCI total daily % dose of DCI // Refills s
Initiate DOJOLVI over a period of For patients swit Discontinue use dosage by appro The total daily d • Caloric value c • Round the tot. • Divide the tot. DOJOLVI Prescription (Titration) Prescription Din DOJOLVI Prescription Din (Maintenance) How Supplied: D Please see full Pl authorize PSP to	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting oximately 5% DCI every 2 to 3 da <b>losage (mL) of DOJOLVI is dete</b> of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w al daily dosage into at least 4 ap <b>Initial Total Daily Dose (mL)</b> Rounded to Nearest Whole Number <b>rections</b> Target Total Daily Dose ( Rounded to Nearest Whole Number DOJOLVI (triheptanoin) oral liquid roduct Monograph atwww.ultrate be my designated agent to for- cription drug order. The patient	In triglyceride (MCT) produ iximately 10% DCI divided i pDOJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin hole number proximately equal individu $\dot{+}$ Doses/Day = (at least 4) $\dot{+}$ (at is supplied in 500 mL bott genyx.com/canada/medici ward this prescription by fa	ct         nto at least 4 times per         // at the last tolerated of up to 35% DCl is ach         of up to 35% DCl is ach         og calculation:         al doses         Initial mL per Dose	r day and increase to the recommended dosage of MCT divided into at least 4 tim nieved.  Total Daily Dose (mL)=  Patients D until reaching targetmL d Use the Prescription Directions field belo describe alternate desired dosing proto  Days Supply  Donograph-CANADA/ for complete dosa bivery, to the pharmacy chosen by the alternate dosine by the alternate do	total daily dosage of up hes per day. Increase the CI (kcal) x Target 8.3 kcal of DOJOLI day(s) Days Supply Refill age and administration ir	to 35% DCI total daily % dose of DC/ // Refills s
Initiate DOJOLVI over a period of For patients swil Discontinue use dosage by appro The total daily d • Caloric value c • Round the tot. • Divide the tot. • Divide the tot. • Divide the tot. • DOJOLVI Prescription (Titration) Prescription Din DOJOLVI Prescription Din (Maintenance) How Supplied: E Please see full Pl authorize PSP to the original pres	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting oximately 5% DCI every 2 to 3 da <b>losage (mL) of DOJOLVI is dete</b> of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w al daily dosage into at least 4 ap <b>Initial Total Daily Dose (mL)</b> Rounded to Nearest Whole Number <b>rections</b> Target Total Daily Dose ( Rounded to Nearest Whole Number 00JOLVI (triheptanoin) oral liquid roduct Monograph atwww.ultrage b be my designated agent to for cription drug order. The patient	In triglyceride (MCT) produ iximately 10% DCI divided i pOJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin hole number proximately equal individu $\dot{=}$ Doses/Day = (at least 4) mL) $\dot{=}$ (at lis supplied in 500 mL bott genyx.com/canada/medici ward this prescription by fa 's chosen pharmacy is the compared to the factor is chosen pharmacy is the compared to the factor (at least 4)	ct         nto at least 4 times per least 4 times 4 times table 4 times table 4 times table 4 times 4 times table 4 times table 4 times 4 times table 4 times	r day and increase to the recommended dosage of MCT divided into at least 4 tim nieved.  Total Daily Dose (mL)= Patients D until reaching targetmL d Use the Prescription Directions field belo describe alternate desired dosing proto  Days Supply  Days Supply  Date	total daily dosage of up hes per day. Increase the CI (kcal) x Target 8.3 kcal of DOJOLI day(s) Days Supply Refill age and administration ir	to 35% DCI total daily % dose of DC/ // Refills s
Initiate DOJOLVI over a period of For patients swill Discontinue use dosage by appro The total daily d • Caloric value c • Round the tot. • Divide the tot. • Divide the tot. • DOJOLVI Prescription (Titration) Prescription Din DOJOLVI Prescription Din (Maintenance) How Supplied: D Please see full Pl authorize PSP to the original pres Prescriber Signal	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting oximately 5% DCI every 2 to 3 da losage (mL) of DOJOLVI is deter of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w al daily dosage into at least 4 ap Initial Total Daily Dose (mL) Rounded to Nearest Whole Number Prections Target Total Daily Dose ( Rounded to Nearest Whole Number DOJOLVI (triheptanoin) oral liquid roduct Monograph atwww.ultrae to be my designated agent to for cription drug order. The patient s. (g, Allergies)	In triglyceride (MCT) produ iximately 10% DCI divided i pDJOLVI. Initiate DOJOLV ays until the target dosage ermined using the followin hole number proximately equal individu (at least 4) mL) ±(at least 4) mL) is supplied in 500 mL bott genyx.com/canada/medici ward this prescription by fa 's chosen pharmacy is the compared to the compared totte to the compared to the compared to the compared totte to the c	ct         nto at least 4 times per         // at the last tolerated of up to 35% DCl is ach         ig calculation:         al doses         Initial mL         per Dose	r day and increase to the recommended dosage of MCT divided into at least 4 tim ieved.  Total Daily Dose (mL)= Patients D until reaching targetmL d Use the Prescription Directions field belo describe alternate desired dosing proto Days Supply Donograph-CANADA/ for complete dosa divery, to the pharmacy chosen by the a and there are no others. Date	total daily dosage of up nes per day. Increase the Cl (kcal) × Target 8.3 kcal of DOJOLI day(s) Days Supply w to cols Refill age and administration ir bove named. This prescr	to 35% DCI total daily % dose of DCI // Refills s
Initiate DOJOLVI over a period of For patients swill Discontinue use dosage by appro The total daily d • Caloric value c • Round the tot. • Divide the tot. • DoJOLVI Prescription (Titration) Prescription Dir DOJOLVI Prescription Dir DOJOLVI Prescription Dir DOJOLVI Prescription C (Maintenance) How Supplied: D Please see full Pr authorize PSP to the original pres Prescriber Signat Special Instruction Special Precaution The prescriber a	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting oximately 5% DCI every 2 to 3 da losage (mL) of DOJOLVI is detect of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest w al daily dosage into at least 4 ap Initial Total Daily Dose (mL) Rounded to Nearest Whole Number Prections Target Total Daily Dose ( Rounded to Nearest Whole Number DOJOLVI (triheptanoin) oral liquid roduct Monograph atww.ultrage be my designated agent to for- cription drug order. The patient turens nsns	In triglyceride (MCT) produ in triglyceride (MCT) produ- in triglyceride (MCT) produ- in triglyceride (MCT) produ- in trigly (MCT) produ- and the product of the produ	ct         nto at least 4 times per         // at the last tolerated of up to 35% DCl is ach         ig calculation:         al doses         Initial mL per Dose	r day and increase to the recommended dosage of MCT divided into at least 4 tim nieved.  Total Daily Dose (mL)= Patients D until reaching targetmL d Use the Prescription Directions field belo describe alternate desired dosing proto  Days Supply  Days Supply  Date	total daily dosage of up hes per day. Increase the CI (kcal) x Target 8.3 kcal of DOJOLI day(s) ose w to cols Days Supply Refill age and administration ir bove named. This prescr	to 35% DCI total daily % dose of DC/ // Refills s
Initiate DOJOLVI over a period of For patients swift Discontinue use dosage by appro The total daily d • Caloric value c • Round the tot. • Divide the tot.	at a total daily dosage of appro 2 to 3 weeks. tching from another MCT produc of MCT products before starting oximately 5% DCI every 2 to 3 da losage (mL) of DOJOLVI is detect of DOJOLVI = 8.3 kcal/mL al daily dosage to the nearest wal daily dosage into at least 4 ap Initial Total Daily Dose (mL) Rounded to Nearest Whole Number Prections Target Total Daily Dose (C Rounded to Nearest Whole Number DOJOLVI (triheptanoin) oral liquid roduct Monograph atww.ultrage be my designated agent to for cription drug order. The patient ture ns (eg, Allergies) ussumes responsibility for mon ME:	In triglyceride (MCT) produ in triglyceride (MCT) produ- in triglyceride (MCT) produ- in triglyceride (MCT) produ- in trigly (MCT) produ- and the product of the produ	ct         nto at least 4 times per         // at the last tolerated of up to 35% DCl is ach         of up to 35% DCl is ach         og calculation:         al doses         Initial mL per Dose	r day and increase to the recommended dosage of MCT divided into at least 4 tim ieved.  Total Daily Dose (mL) = Patients D Increase bymL everyd Use the Prescription Directions field belo describe alternate desired dosing proto  Days Supply  Days Supply  Date  nsibility for notifying UltraCare of any  Darticipate in relevant Ultragenyx marked	total daily dosage of up hes per day. Increase the CI (kcal) x Target 8.3 kcal of DOJOLI day(s) ose w to cols Days Supply Refill age and administration ir bove named. This prescr	to 35% DCI total daily % dose of DC/ // Refills s formation. I iption represen

© 2024 Ultragenyx Pharmaceutical Inc. All rights reserved. DOJOLVI is a registered trademark of Ultragenyx Pharmaceutical, Inc. MRCP-UX007-00309 09/2024